

# Long-term storage of feces at $-80^{\circ}\text{C}$ versus $-20^{\circ}\text{C}$ is negligible for 16S rRNA amplicon profiling of the equine bacterial microbiome

Stefan Gavriiliuc, Mason R. Stothart, Astrid Henry and Jocelyn Poissant

Faculty of Veterinary Medicine, University of Calgary, Calgary, AB, Canada

## ABSTRACT

The development of next-generation sequencing technologies has spurred a surge of research on bacterial microbiome diversity and function. But despite the rapid growth of the field, many uncertainties remain regarding the impact of differing methodologies on downstream results. Sample storage temperature is conventionally thought to be among the most important factors for ensuring reproducibility across marker gene studies, but to date much of the research on this topic has focused on short-term storage in the context of clinical applications. Consequently, it has remained unclear if storage at  $-80^{\circ}\text{C}$ , widely viewed as the gold standard for long-term archival of feces, is truly required for maintaining sample integrity in amplicon-based studies. A better understanding of the impacts of long-term storage conditions is important given the substantial cost and limited availability of ultra-low temperature freezers. To this end, we compared bacterial microbiome profiles inferred from 16S V3–V4 amplicon sequencing for paired fecal samples obtained from a feral horse population from Sable Island, Nova Scotia, Canada, stored at either  $-80^{\circ}\text{C}$  or  $-20^{\circ}\text{C}$  for 4 years. We found that storage temperature did not significantly affect alpha diversity measures, including amplicon sequence variant (ASV) richness and evenness, and abundance of rare sequence variants, nor presence/absence, relative abundances and phylogenetic diversity weighted measures of beta diversity. These results indicate that storage of equine feces at  $-20^{\circ}\text{C}$  for periods ranging from a few months to a few years is equivalent to storage at  $-80^{\circ}\text{C}$  for amplicon-based microbiome studies, adding to accumulating evidence indicating that standard domestic freezers are both economical and effective for microbiome research.

Submitted 4 November 2020

Accepted 4 January 2021

Published 9 March 2021

Corresponding authors

Stefan Gavriiliuc,  
stefan.gavriiliuc@ucalgary.ca

Jocelyn Poissant,  
jocelyn.poissant@ucalgary.ca

Academic editor

Elliot Lefkowitz

Additional Information and  
Declarations can be found on  
page 11

DOI [10.7717/peerj.10837](https://doi.org/10.7717/peerj.10837)

© Copyright

2021 Gavriiliuc et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Biodiversity, Bioinformatics, Ecology, Microbiology, Veterinary Medicine

**Keywords** Microbiome, 16S, DNA metabarcoding, Long-term storage, *Equus ferus caballus*, Protocol, Sequencing, Horse, Amplicon, Equine

## INTRODUCTION

Targeted marker gene sequencing allows for the characterization of biological communities at high-resolution (*Cristescu, 2014*). In particular, quantifying the resident bacteria of the gastrointestinal tract, or the gut microbiome, using deep sequencing of 16S rRNA amplicons has furthered our understanding of host physiology and disease

etiology, and improved clinical diagnostics and treatment of gut dysbiosis (Kim *et al.*, 2017; Vandeputte *et al.*, 2017). As the cost of Next-Generation Sequencing (NGS) experiments continues to fall, marker gene surveys have become increasingly commonplace. However, the findings of different studies are often not readily comparable or generalizable because of a lack of standardized methodology (Vandeputte *et al.*, 2017). For example, the use of different DNA extraction kits and amplification methodologies have been observed to introduce technical variation comparable to that of biological variation in metagenomic and amplicon-based experiments (Costea *et al.*, 2017; Sinha *et al.*, 2017). While these effects are substantial and well-documented, the spread of marker gene surveys has brought with it a renewed interest in optimizing and finding alternative sample storage solutions, which are also known as large sources of technical variation (Kim *et al.*, 2017).

Short-term processing and storage of fecal samples for microbiome research has received considerable attention due to their importance in clinical applications (Table 1). Where immediate extraction is not possible, the conventionally accepted gold standard for short-term storage is rapid freezing at  $-80^{\circ}\text{C}$  without buffer (Vandeputte *et al.*, 2017). While the process of freezing samples is still under contention (Bahl, Bergström & Licht, 2012; Fouhy *et al.*, 2015; Metzler-Zebeli *et al.*, 2016), it remains by far the most common approach for preventing rapid changes in the abundance of specific taxa under ambient conditions following sampling. This phenomenon, often referred to as microbial 'blooms', has been well documented. For example, a meta-analysis has shown that human rectal swabs stored at room temperature experience characteristic growths of *Gammaproteobacteria* (Amir *et al.*, 2017). A similar increase in the abundance of *proteobacteria* was also seen in the microbiota of tadpoles (*Nanorana parker*) gut samples left thawing at ambient temperature for 12 h (Anslan *et al.*, 2019), while equine feces collected 12 h post-defecation showed sharp increases of families from the *Firmicutes* phylum (Beckers, Schulz & Childers, 2017). Though the effect of sample storage at ambient temperature has been well-documented across study systems, it has remained unclear whether freezing at  $-80^{\circ}\text{C}$  is overly conservative. Notably, an increasing number of studies have claimed or demonstrated that storage at  $-20^{\circ}\text{C}$  is generally sufficient over the short-term (Song *et al.*, 2016; Blekhman *et al.*, 2016; Panek *et al.*, 2018; Wang *et al.*, 2018). Overall, the consensus emerging from these studies is that (any) differences in microbiome profiles caused by variation in freezing temperature are typically negligible when compared to biological patterns of interest (Lauber *et al.*, 2010; Carroll *et al.*, 2012; Dominianni *et al.*, 2014; Blekhman *et al.*, 2016; Bundgaard-Nielsen, Hagström & Sørensen, 2018) and that storage at  $-80^{\circ}\text{C}$  may therefore be unnecessary for storage periods less than a few weeks or months.

While the impacts of short-term storage temperature on microbiome profiles are relatively well understood, less is known of the impacts of storage conditions over long periods of time. Delineating short-term from long-term storage is necessarily subjective, but a gap exists in the literature for storage periods greater than a few weeks or months (Table 1). A notable exception is the study of Kia *et al.* (2016) which showed that the microbiota from freeze-dried human fecal samples stored for 14 years at  $-20^{\circ}\text{C}$

**Table 1** Summary of studies having investigated the effect of freezing storage temperature ( $-20\text{ }^{\circ}\text{C}$  or  $-80\text{ }^{\circ}\text{C}$ ) on microbiome profiling using 16S amplicon sequencing. Selected studies have compared samples stored at different freezing temperature without buffer. Studies are ordered by storage duration.

Duration	Temp ( $^{\circ}\text{C}$ )	Control	Sample	16S Region	Diversity measure (statistical test/method)	Conclusions	References
~1 day	$-20\text{ }^{\circ}\text{C}$ + 1–2 freeze thaw cycles, then $-80\text{ }^{\circ}\text{C}$	$-80\text{ }^{\circ}\text{C}$	Human stool and rectal swab	V4	$\alpha$ : Richness (Kruskal–Wallis) $\beta$ : Yue & Clayton dissimilarity index (PCoA, Wilcoxon)	$\alpha$ : No significant effect $\beta$ : Samples clustered by host (not storage method), no difference in $\theta_{\text{YC}}$ between storage temperatures	<a href="#">Bassis et al. (2017)</a>
1 or 3 days	$-20\text{ }^{\circ}\text{C}$	$-80\text{ }^{\circ}\text{C}$	Human stool	V4	$\alpha$ : Richness, Shannon (ANOVA & Tukey post-hoc/Kruskal-Wallis & Dunn's post-hoc) $\beta$ : Hellinger distances (PCA) OTU – 25 most abundant taxa	$\alpha$ : No significant effect $\beta$ : Samples clustered by host rather than storage method OTU: No significant effect	<a href="#">Bundgaard-Nielsen, Hagström &amp; Sørensen (2018)</a>
1 week	$-20\text{ }^{\circ}\text{C}$	$-80\text{ }^{\circ}\text{C}$	Human stool	V1–V3	$\alpha$ : Richness, Chao1, Shannon, Phylogenetic Diversity whole tree (Wilcoxon signed rank) $\beta$ : Weighted & unweighted UniFrac, Bray-Curtis (UPGMA & PCoA clustering, Mann–Whitney $U$ test, PERMANOVA) OTU: Pairwise comparisons of specific taxa ( $G$ -test)	$\alpha$ : No significant effect except for Shannon diversity in one sample $\beta$ : Samples clustered by host rather than storage temperature; no significant difference between temperatures OTU: No effect	<a href="#">Tedjo et al. (2015)</a>
3 & 14 days	$-20\text{ }^{\circ}\text{C}$	$-80\text{ }^{\circ}\text{C}$	Human stool & skin, soil	V1–V2	$\alpha$ : Faith's phylogenetic diversity (Kruskal–Wallis) $\beta$ : UniFrac (PERMANOVA, NMDS) OTU: Relative taxon abundance (Kruskal–Wallis)	$\alpha$ : No effect $\beta$ : No effect OTU: minor differences in abundance of specific taxa but no significant effect on overall community composition	<a href="#">Lauber et al. (2010)</a>
4 weeks	$-20\text{ }^{\circ}\text{C}$ , $-80\text{ }^{\circ}\text{C}$	Fresh	Human pharyngeal	V1–V3	$\alpha$ : Richness (Dirichlet multinomial models) $\beta$ : Weighted UniFrac (PCoA, Dirichlet multinomial models) OTU: Relative taxon abundance (metagenomeSeq)	$\alpha$ : No significant effect $\beta$ : No significant effect OTU – No significant effect	<a href="#">Hang et al. (2014)</a>
33 days	$-20\text{ }^{\circ}\text{C}$ , $-80\text{ }^{\circ}\text{C}$	Fresh	Murine stool	V3–V4	$\alpha$ : Richness, Chao1 (two-tailed Student's $t$ -test, non-parametric Mann–Whitney) $\beta$ : Bray-Curtis (PCoA)	$\alpha$ : No significant effect $\beta$ : Samples clustered by host rather than storage temperature)	<a href="#">Jenkins et al. (2018)</a>

(Continued)

Table 1 (continued)

Duration	Temp (°C)	Control	Sample	16S Region	Diversity measure (statistical test/method)	Conclusions	References
5 weeks total	-20 °C (1 week) then -80 °C (4 weeks)	Fresh	Human mid-vaginal swab	V1-V2	$\beta$ : Jensen-Shannon divergence, relative entropy, Euclidean distance, Bray-Curtis (Kolmogorov-Smirnov)	$\beta$ : No difference between storage temperatures	<a href="#">Bai et al. (2012)</a>
Up to 8 weeks	-20 °C, -80 °C	Fresh	Spider monkey stool	V4	$\alpha$ : Shannon, Simpson (LSMEANS) $\beta$ : Weighted & unweighted UniFrac (PERMANOVA, PCoA, distance boxplot analysis) OTU: Mean frequencies (Kruskal-Wallis, decision trees)	$\alpha$ : No significant effect $\beta$ : Fresh samples clustered with frozen samples, frozen sample not significantly different from fresh samples OTU: Frozen samples misassigned to each other	<a href="#">Hale et al. (2015)</a>
3 months	-20 °C, -80 °C	Fresh	Porcine fecal samples	Taxa specific	OTU: Relative and absolute abundances of specific groups (linear discriminant analysis)	OTU: Frozen samples most similar to each other	<a href="#">Metzler-Zebeli et al. (2016)</a>
14 years	Freeze-dried then -20 °C	Matched cohort stored at -80 °C	Human stool	V3-V4	$\beta$ : Bray-Curtis, weighted & unweighted UniFrac (ANOSIM, NMDS)	$\beta$ : Significant difference in Bray-Curtis dissimilarities but not UniFrac distances, clustering overlap between -20 °C and matched cohort	<a href="#">Kia et al. (2016)</a>

were similar to those of samples stored at -80 °C in a cohort matched for age, sex, BMI and pre-existing health conditions ([McDonald et al., 2018](#)). However, a significant difference was nonetheless observed between groups when considering Bray-Curtis dissimilarities, and it is unclear if similar results would be obtained without the use of freeze-drying which is an uncommon practice in microbiome studies.

In the absence of relevant research, -80 °C has been widely adopted as the gold standard for long-term storage of biological samples intended for microbiome analyses including feces ([Hale et al., 2015](#); [Jenkins et al., 2018](#); [Shaw et al., 2016](#)), necessitating the use of ultra-low temperature freezers. Compared to regular -20 °C freezers, ultra-low temperature freezers incur higher capital and energy costs, have higher carbon footprints, and lower durability ([Gumapas & Simons, 2013](#)). In addition to constraints imposed by those with limited funding and resources ([Vogtmann et al., 2017](#); [Carruthers et al., 2019](#)), immediate or prompt transfer to -80 °C freezers is also often logistically impractical, especially for studies conducted in remote areas ([Song et al., 2016](#); [Blekhman et al., 2016](#)). Ultimately, meeting current gold standards often impose significant financial and logistical burdens on researchers, and failure to meet these standards creates uncertainty regarding sample integrity and the validity and comparability of results.

Given the need to validate storage protocols currently in use by our group and others for projects where access to  $-80^{\circ}\text{C}$  storage is limited or impossible, we evaluated the effect of long-term freezing storage temperature on fecal samples collected as part of a long-term study of feral horses on Sable Island, Nova Scotia, Canada (see [Gold et al., 2019](#) and [Regan et al., 2020](#) for details about this study). Specifically, we tested whether storing aliquots from the same fecal sample at either  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for 4 years affected 16S rRNA amplicon characterization of the equine bacterial microbiome. This study is among the firsts to explicitly test the for impacts of long-term freezing temperature on 16S microbiome profiling, and will contribute to protocol development and evaluation in microbiome research.

## MATERIALS AND METHODS

### Field samples

The fecal samples used in this study were obtained from the population of feral horses on Sable Island, Nova Scotia, Canada ( $43^{\circ}55'\text{N}$ ;  $-60^{\circ}00'\text{W}$ ). We collected the surface portion (not in contact with the ground) of eight freshly voided fecal samples using nitrile gloves which were then inverted and sealed. Samples were kept on ice in the field for a maximum of 7 h before being sub-sampled into 1–2 mL cryovials on the same day. Paired samples (aliquots) were then either stored immediately in a liquid nitrogen dewar and subsequently transferred to a  $-80^{\circ}\text{C}$  freezer or stored immediately in  $-20^{\circ}\text{C}$  chest freezers for 4 years. Sample collection and subsequent laboratory analyses were performed under Parks Canada Agency Research and Collections Permit SINP-2013-2014, University of Saskatchewan Animal Care Protocol 20090032, and University of Calgary Animal Care Protocol AC18-0078.

### DNA extraction and sequencing

We extracted DNA from 0.2 g sub-samples of feces using a single Qiagen Powersoil kit (Qiagen, Hilden, Germany) following the manufacturer default recommendations. The protocol followed Qiagen's Kit Handbook with homogenization performed at full speed for 10 min using the 2 mL bead beating tubes provided with the kit (0.7 mm Dry Garnet) and a Vortex-Genie 2 fitted with Qiagen's Vortex Adapter (Cat. No. 13000-V1-24). All samples were vortexed simultaneously with paired samples distributed randomly across the Vortex Adapter. DNA extracts were quantified using the Qubit dsDNA BR Assay Kit and standardized to 20 ng/uL with molecular grade water prior to polymerase chain reaction (PCR) amplification. PCR and sequencing was outsourced to the University of Calgary Centre for Health Genomics and Informatics. Library preparation, including PCR program and conditions, was done as specified in the Illumina 16S Metagenomic Sequencing Library Preparation protocol Part # 15044223 Rev. B. Briefly, PCR was conducted on the V3–V4 region of the 16S rRNA gene using the 341F/805R Primers. Sequencing was conducted on an Illumina MiSeq platform using v3 chemistry (600-cycle) to generate  $2 \times 300$  base pair (bp) paired-end reads using dual 8 bp indexing, a loading concentration of 4 pM and 25% phiX spike-ins. Lastly, we used 2

blanks containing only water and a no-template control (NTC) as negative controls to assess for contamination from extraction and sequencing.

## Bioinformatics


Raw reads were demultiplexed and indices removed during the MiSeq post-run processing. Cutadapt v2.8 (Martin, 2011) was used to remove sequencing primers in paired-end mode with the number of searches for a primer in a sequence raised to  $n = 5$ . Sequences with ambiguous nucleotides were removed using the *filterAndTrim* function from the DADA2 package (Callahan et al., 2016). DADA2 v1.16 with the corresponding tutorial was used to process primerless sequences and extract amplicon sequence variants (ASVs). Quality filtering was conducted with truncation lengths set to 250 and 200 for forward and reverse reads, respectively, while the maximum number of expected errors were left unchanged. The number of rounds to convergence was increased from 10 to 15 for constructing empirical sequencing error models. ASVs were extracted using the core *dada* algorithm with the option for pooling sequences set to TRUE. Forward and reverse reads were subsequently merged using default parameters with *mergePairs*, and chimeras were removed using the *consensus* method. ASVs represented by a single copy in the final ASV table (singletons) were excluded.

Taxonomy was assigned using the DADA2 implementation of the RDP classifier algorithm, *assignTaxonomy*, with the Silva 138 reference taxonomy database and an 80% confidence threshold. For phylogenetic-based beta diversity analyses, we aligned all ASVs using *AlignSeqs* from the DECIPHER package v2.16.1 (Wright, 2015) and created a neighbor-joining tree using the *NJ* command from the *phangorn* package v2.5.5 (Schliep, 2011) using default parameters. Finally, we used the *decontam* v1.8.0 package (Davis et al., 2018) to remove contaminant ASVs using the prevalence method and the default threshold for detection.

## Statistics

Statistical analysis was conducted with the *vegan* v2.5-6 (Oksanen et al., 2019) and *phyloseq* v1.32.0 (McMurdie & Holmes, 2013) packages unless otherwise noted. Prior to statistical analysis, samples were rarefied to the minimum sequencing depth (45,832) among all samples using *rarefy\_even\_depth()*. We quantified alpha diversity using observed richness, Shannon and Chao1 diversity indices for rarefied ASV counts. Observed richness and the Chao1 index were normally distributed and compared between temperature treatments using paired two-sided *t*-tests, while Shannon indices were non-normally distributed and therefore compared using a Wilcoxon signed-rank test.

We calculated distance matrices using two phylogeny-independent (Euclidean and Jaccard distances; Jaccard, 1912) and phylogeny-informed (weighted and unweighted UniFrac distances; Lozupone & Knight, 2005) beta diversity distance measures. Notably, to obtain Euclidean distances that satisfy the compositionality constraint of amplicon sequence data (Luz Calle, 2019), we applied a centered-log ratio transformation to unrarefied counts using the *microbiome* package v1.10.0 (Lahti & Sudarshan, 2017). Patterns of beta diversity were visualized using ordinations, and to test for the influence of



**Figure 1** Relative abundance of bacterial families identified in fecal samples collected from feral horses on Sable Island, Nova Scotia, Canada, using 16S rRNA sequencing. Samples from eight horses were divided into aliquots, stored at both  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for 4 years, and characterized using 16S amplicon sequencing. Taxonomy was assigned to amplicon sequence variants (ASVs), and abundances were converted to proportional values then aggregated to the family level. Families with low abundances within a sample ( $<1\%$ ) were pooled for clarity. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04\_img.jpg\) DOI: 10.7717/peerj.10837/fig-1](https://doi.org/10.7717/peerj.10837/fig-1)


Sample ID and storage temperature on diversity we ran Permutational Multivariate Analyses of Variance (PERMANOVA) including both effects using the *adonis2* function from *vegan* on the four aforementioned distance matrices at the level of ASVs.

The *betadisper* function from *vegan* was also used to test whether the variance in compositional abundances differed between storage temperatures.

## RESULTS

### Community composition

Sequencing generated a total of 1,615,043 reads, with a mean of 100,940 reads per sample (87,049–124,423). After sequence processing including quality filtering, denoising, merging and chimera removal, an average of 52,600 reads remained per sample (45,827–64,386). A total of 4427 ASVs were identified, of which 126 were singletons and a further 33 were identified as contaminants. After removing singletons and contaminants, an average of 71.0%, 88.0% and 97.7% of sequences within a sample were assigned to the genus, family and phylum levels with at least 80% confidence, respectively. Bacterial communities were dominated by the *Firmicutes* (average proportion 40.8%;



**Figure 2** Alpha diversity (richness (A), Chao1 (B) and Shannon indices (C)) of 16S V3–V4 Amplicon Sequence Variants (ASV) for paired equine fecal samples (aliquots) stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for 4 years. Samples were collected from separate individuals on Sable Island, Nova Scotia, Canada. Paired samples (aliquots) are denoted by color. [Full-size !\[\]\(5fd6ef84f97f42d7f8b34275f1b65312\_img.jpg\) DOI: 10.7717/peerj.10837/fig-2](https://doi.org/10.7717/peerj.10837/fig-2)

32.1–53.2%) and *Bacteroidota* phyla (average proportion 35.5%; 22.3–41.4%, Fig. S1). A total of 89 unique families were observed, and 65 of these were present at proportional abundances of 1% or less within a sample (Fig. 1). At the genus level, 138 unique genera were identified of which 114 were present at proportional abundances of 1% or less within a sample (Fig. S2).


### Alpha diversity comparisons

The number of unique ASVs per sample ranged from 1,565 to 2,095 with a mean ( $\pm 1$  standard deviation) of  $1,797 \pm 172$  (Fig. 2). Chao1 estimates ranged from 1,677 to 2,305 with a mean of  $1,935 \pm 200$ , and Shannon values ranged from 5.79 to 6.44 with a mean of  $6.00 \pm 0.17$  (Fig. 2). Observed ASV richness ( $p = 0.39$ ), Chao1 indices ( $p = 0.64$ ) and Shannon diversity ( $p = 0.11$ ) did not differ significantly between samples stored at different temperatures.

### Beta diversity comparisons

Aliquots from the same sample generally yielded consistent results, as apparent in NMDS ordinations (Fig. 3). The proportion of multivariate variance explained by sample ID was large (79–88%) and statistically significant ( $p < 0.001$ ) for all distance measures





**Figure 3** Non-metric multidimensional scaling (NMDS) of equine gut bacterial communities inferred from fecal samples stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$ . Ordinations were conducted on (A) Euclidean (stress = 0.08), (B) Jaccard (stress = 0.10), (C) Unweighted UniFrac (stress = 0.10) and (D) weighted UniFrac (stress = 0.14) distance matrices from Amplicon Sequencing Variant (ASV) count data. Paired samples share the same color and temperature is denoted by circles ( $-20^{\circ}\text{C}$ ) and triangles ( $-80^{\circ}\text{C}$ ). Red and blue ellipses represent 95% confidence intervals for samples stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$ , respectively. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674\_img.jpg\) DOI: 10.7717/peerj.10837/fig-3](https://doi.org/10.7717/peerj.10837/fig-3)

considered (Table S1). In contrast, storage temperature explained a comparatively small (2–3%) and non-significant amount of multivariate variation across distances. Similarly, no difference in beta dispersion was observed between temperature storage groups indicating homogeneity of variance in the community structure between treatment groups (Table S1).

## DISCUSSION

Immediate extraction of DNA for amplicon sequencing experiments is preferable but many study systems and clinical applications require the storage of samples by way of freezing. We aimed to determine if equine fecal samples stored at  $-20^{\circ}\text{C}$  for multiple years could recapitulate the bacterial community structure inferred from samples stored at  $-80^{\circ}\text{C}$ . Samples stored at different temperatures did not differ in ASVs richness, Chao1 indices, or evenness (Shannon). Furthermore, variation in community composition among samples was largely driven by sample (individual) ID with no detectable impact of storage temperature. In short, our study demonstrates that archiving equine feces at either  $-20^{\circ}\text{C}$  or  $-80^{\circ}\text{C}$  for multiple years has negligible impacts of 16S rRNA microbiome profiling.

Our findings mirror those obtained in similar studies that have looked at the impact of storage freezing temperature on microbiome profiling (Table 1). For example, *Jenkins et al. (2018)* noted that murine fecal samples stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for 33 days did not differ in terms of richness or the Chao1 estimator. Microbiome profiles from spider monkey feces stored at  $-20^{\circ}\text{C}$  for 8 weeks closely resembled those stored at  $-80^{\circ}\text{C}$  in composition and diversity (*Hale et al., 2015*). Using pyrosequencing reads, *Hang et al. (2014)* did not observe differences between oropharyngeal swabs at these temperatures using weighted UniFrac distances.

Though we did not compare frozen samples to freshly extracted samples, previous studies have shown that long-term storage at  $-20^{\circ}\text{C}$  generally gives rise to microbiota profiles similar to those that have been immediately extracted (*Hale et al., 2015; Kia et al., 2016*). In addition, limited differences were observed between fresh fecal samples and those stored at  $-80^{\circ}\text{C}$  in RNAlater for 5 years (*Tap et al., 2019*). Together with our findings, these suggest that archiving fecal samples for long periods of time at  $-20^{\circ}\text{C}$  likely yields uncompromised microbiome profiles, even when compared to DNA extracted from fresh samples. Studies comparing results from fresh fecal samples with those of samples archived at  $-20^{\circ}\text{C}$  for multiple aliquots would be valuable.

Based on this study and other accumulating evidence, we suggest the use of  $-20^{\circ}\text{C}$  as a cost-effective storage temperature to maintain microbiome integrity over multiple years comparable to that of samples stored at  $-80^{\circ}\text{C}$ , at least for targeted sequencing of short amplicons. However, it remains to be seen if this is generalizable to longer time periods such as decades, other sample types or host organisms, or non-amplicon based sequencing methods such as shotgun metagenomics and third generation sequencing. It is also possible that larger sample sizes would lead to the detection of subtle variation, if present. We also note that our study used manual defrost freezers, and that long-term archival of samples in self-defrosting freezers experiencing frequent temperature fluctuations may yield different outcomes.

In addition to general relevance, our findings are especially pertinent to clinicians and researchers conducting field or community-based studies where access to  $-80^{\circ}\text{C}$  storage is limited or absent. Wider adoption and acceptance of storage at  $-20^{\circ}\text{C}$  for amplicon-based microbiome research should greatly increase total storage space, encourage the re-use of samples, and promote longitudinal sampling (*Björk et al., 2019*). Increased reliance on  $-20^{\circ}\text{C}$  freezers, at least for storage up to a few years, may also help reduce drop-out rates in microbiome studies by alleviating the logistical burden of rapidly transferring samples to  $-80^{\circ}\text{C}$  (*Vandeputte et al., 2017*). In terms of standardizing study design, DNA extraction and amplification protocols may be of greater importance for resource allocation, which are likely larger sources of technical variation (*Costea et al., 2017; Sinha et al., 2017*) compared to long-term storage temperature.

## CONCLUSION

We demonstrated that technical replicates of equine fecal samples stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for multiple years yield similar bacterial microbiome results in terms of alpha

and beta diversity. Our findings add to mounting evidence indicating that standard domestic freezers are both economical and effective for microbiome research.

## ACKNOWLEDGEMENTS

We thank the students, research assistants and volunteers who contributed to sample collection, Philip McLoughlin for coordinating field activities and transport of samples, and Matthew Workentine for bioinformatic support. In-kind and logistical support was provided by Fisheries and Oceans Canada (DFO), Environment Canada, Parks Canada Agency, Maritime Air Charters Limited (Sable Aviation), and Sable Island Station (Meteorological Service of Canada).

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the University of Calgary, the Margaret Gunn Endowment for Animal Research, the L. David Dubé and Heather Ryan Veterinary Health and Research Fund, and Natural Sciences and Engineering Research Council of Canada Discovery Grants Nos. 2019-04388 to Jocelyn Poissant and 2016-06459 to collaborator Philip D. McLoughlin. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

University of Calgary.

Margaret Gunn Endowment.

L. David Dubé and Heather Ryan Veterinary Health and Research Fund.

Natural Sciences and Engineering Research Council of Canada Discovery: 2019-04388 and 2016-06459.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Stefan Gavriiliuc conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Mason R. Stothart conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Astrid Henry conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Jocelyn Poissant conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

## Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Field sampling was approved by the University of Saskatchewan Animal Care Committee (Protocol 20090032) and the University of Calgary Animal Care Committee (Protocol AC18-0078).

## Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Sample collection and subsequent laboratory analyses were performed under Parks Canada Agency Research and Collections Permit SINP-2013-2014.

## Data Availability

The following information was supplied regarding data availability:

Raw 16S sequences are available at NCBI, BioProject: [PRJNA674675](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA674675) (SRX9507884–SRX9507891, SRX9440373, SRX9440395, SRX9440405, SRX9440422, SRX9440423, SRX9440428, SRX9440439, SRX9440443, SRX9517110, SRX9517111, SRX9732812).

Raw sequences and code are available as [Supplemental Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10837#supplemental-information>.

## REFERENCES

- Amir A, McDonald D, Navas-Molina JA, Debelius J, Morton JT, Hyde E, Robbins- Pianka A, Knight R. 2017. Correcting for microbial blooms in fecal samples during room-temperature shipping. *mSystems* 2(2):e00199-16 DOI [10.1128/msystems.00199-16](https://doi.org/10.1128/msystems.00199-16).
- Anslan S, Li H, Künzel S, Vences M. 2019. Microbiomes from feces vs. gut in aquatic vertebrates: distinct community compositions between substrates and preservation methods. *BioRxiv* 651612:e01294 DOI [10.1101/651612](https://doi.org/10.1101/651612).
- Bahl MI, Bergström A, Licht TR. 2012. Freezing fecal samples prior to DNA extraction affects the firmicutes to bacteroidetes ratio determined by downstream quantitative PCR analysis. *FEMS Microbiology Letters* 329(2):193–197 DOI [10.1111/j.1574-6968.2012.02523.x](https://doi.org/10.1111/j.1574-6968.2012.02523.x).
- Bai G, Gajer P, Nandy M, Ma B, Yang H, Sakamoto J, Blanchard MH, Ravel J, Brotman RM. 2012. Comparison of storage conditions for human vaginal microbiome studies. *PLOS ONE* 7(5):e36934 DOI [10.1371/journal.pone.0036934](https://doi.org/10.1371/journal.pone.0036934).
- Bassis CM, Moore NM, Lolans K, Seekatz AM, Weinstein RA, Young VB, Hayden MK. 2017. Comparison of stool versus rectal swab samples and storage conditions on bacterial community profiles. *BMC Microbiology* 17(1):1–7 DOI [10.1186/s12866-017-0983-9](https://doi.org/10.1186/s12866-017-0983-9).
- Beckers KF, Schulz CJ, Childers GW. 2017. Rapid regrowth and detection of microbial contaminants in equine fecal microbiome samples. *PLOS ONE* 12(11):e0187044 DOI [10.1371/journal.pone.0187044](https://doi.org/10.1371/journal.pone.0187044).

- Björk JR, Dasari M, Grieneisen L, Archie EA. 2019.** Primate microbiomes over time: longitudinal answers to standing questions in microbiome research. *American Journal of Primatology* **81**(10–11):e22970 DOI [10.1002/ajp.22970](https://doi.org/10.1002/ajp.22970).
- Blekhman R, Tang K, Archie EA, Barreiro LB, Johnson ZP, Wilson ME, Kohn J, Yuan ML, Gesquiere L, Grieneisen LE, Tung J. 2016.** Common methods for fecal sample storage in field studies yield consistent signatures of individual identity in microbiome sequencing data. *Scientific Reports* **6**(1):31519 DOI [10.1038/srep31519](https://doi.org/10.1038/srep31519).
- Bundgaard-Nielsen C, Hagstrøm S, Sørensen S. 2018.** Interpersonal variations in gut microbiota profiles supersedes the effects of differing fecal storage conditions. *Scientific Reports* **8**(1):17367 DOI [10.1038/s41598-018-35843-0](https://doi.org/10.1038/s41598-018-35843-0).
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.** DADA2: high-resolution sample inference from illumina amplicon data. *Nature Methods* **13**(7):581–583 DOI [10.1038/nmeth.3869](https://doi.org/10.1038/nmeth.3869).
- Carroll IM, Ringel-Kulka T, Siddle JP, Klaenhammer TR, Ringel Y. 2012.** Characterization of the fecal microbiota using high-throughput sequencing reveals a stable microbial community during storage. *PLOS ONE* **7**(10):e46953 DOI [10.1371/journal.pone.0046953](https://doi.org/10.1371/journal.pone.0046953).
- Carruthers LV, Moses A, Adriko M, Faust CL, Tukahebwa EM, Hall LJ, Ranford-Cartwright LC, Lamberton PHL. 2019.** The impact of storage conditions on human stool 16S rRNA microbiome composition and diversity. *PeerJ* **7**(1):e8133 DOI [10.7717/peerj.8133](https://doi.org/10.7717/peerj.8133).
- Costea PI, Zeller G, Sunagawa S, Pelletier E, Alberti A, Levenez F, Tramontano M, Driessen M, Hercog R, Jung F-E, Kultima JR, Hayward MR, Coelho LP, Allen-Vercoe E, Bertrand L, Blaut M, Brown JRM, Carton T, Séphanie C-P, Daigneault M, Derrien M, Druesne A, De Vos WM, Finlay BB, Flint HJ, Guarner F, Hattori M, Heilig H, Luna RA, Van Hylckama Vlieg J, Junick J, Klymiuk I, Langella P, Le Chatelier E, Mai V, Manichanh C, Martin JC, Cémentine M, Morita H, O’Toole PW, Céline O, Patil KR, Penders J, Søren P, Pons N, Popova M, Salonen A, Saulnier D, Scott KP, Singh B, Slezak K, Veiga P, Versalovic J, Zhao L, Zoetendal EG, Ehrlich SD, Dore J, Bork P. 2017.** Towards standards for human fecal sample processing in metagenomic studies. *Nature Biotechnology* **35**:1069–1076 DOI [10.1038/nbt.3960](https://doi.org/10.1038/nbt.3960).
- Cristescu ME. 2014.** From barcoding single individuals to metabarcoding biological communities: towards an integrative approach to the study of global biodiversity. *Trends in Ecology and Evolution* **29**(10):566–571 DOI [10.1016/j.tree.2014.08.001](https://doi.org/10.1016/j.tree.2014.08.001).
- Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018.** Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome* **6**(1):226 DOI [10.1186/s40168-018-0605-2](https://doi.org/10.1186/s40168-018-0605-2).
- Dominianni C, Wu J, Hayes RB, Ahn J. 2014.** Comparison of methods for fecal microbiome biospecimen collection. *BMC Microbiology* **14**(1):103 DOI [10.1186/1471-2180-14-103](https://doi.org/10.1186/1471-2180-14-103).
- Fouhy F, Deane J, Rea MC, O’Sullivan Ó, Ross RP, O’Callaghan G, Plant BJ, Stanton C. 2015.** The effects of freezing on faecal microbiota as determined using MiSeq sequencing and culture-based investigations. *PLOS ONE* **10**(3):e0119355 DOI [10.1371/journal.pone.0119355](https://doi.org/10.1371/journal.pone.0119355).
- Gold S, Regan CE, McLoughlin PD, Gilleard JS, Wilson AJ, Poissant J. 2019.** Quantitative genetics of gastrointestinal strongyle burden and associated body condition in feral horses. *International Journal for Parasitology: Parasites and Wildlife* **9**:104–111 DOI [10.1016/j.ijppaw.2019.03.010](https://doi.org/10.1016/j.ijppaw.2019.03.010).
- Gumapas LAM, Simons G. 2013.** Factors affecting the performance, energy consumption, and carbon footprint for ultra low temperature freezers: case study at the National Institutes of

- Health. *World Review of Science, Technology and Sustainable Development* **10(1/2/3)**:129  
DOI [10.1504/WRSTSD.2013.050786](https://doi.org/10.1504/WRSTSD.2013.050786).
- Hale VL, Tan CL, Knight R, Amato KR. 2015. Effect of preservation method on spider monkey (*Ateles geoffroyi*) fecal microbiota over 8 weeks. *Journal of Microbiological Methods* **113**:16–26  
DOI [10.1016/j.mimet.2015.03.021](https://doi.org/10.1016/j.mimet.2015.03.021).
- Hang J, Desai V, Zavaljevski N, Yang Y, Lin X, Satya RV, Martinez LJ, Blaylock JM, Jarman RG, Thomas SJ, Kushner RA. 2014. 16S rRNA gene pyrosequencing of reference and clinical samples and investigation of the temperature stability of microbiome profiles. *Microbiome* **2(1)**:31 DOI [10.1186/2049-2618-2-31](https://doi.org/10.1186/2049-2618-2-31).
- Jaccard P. 1912. The distribution of the flora in the alpine zone. *New Phytologist* **11(2)**:37–50  
DOI [10.1111/j.1469-8137.1912.tb05611.x](https://doi.org/10.1111/j.1469-8137.1912.tb05611.x).
- Jenkins SV, Vang KB, Gies A, Griffin RJ, Jun SR, Nookaew I, Dings RPM. 2018. Sample storage conditions induce post-collection biases in microbiome profiles. *BMC Microbiology* **18(1)**:227  
DOI [10.1186/s12866-018-1359-5](https://doi.org/10.1186/s12866-018-1359-5).
- Kia E, MacKenzie BW, Middleton D, Lau A, Waite DW, Lewis G, Chan YK, Silvestre M, Cooper GJS, Poppitt SD, Taylor MW. 2016. Integrity of the human faecal microbiota following long-term sample storage. *PLOS ONE* **11(10)**:e0163666 DOI [10.1371/journal.pone.0163666](https://doi.org/10.1371/journal.pone.0163666).
- Kim D, Hofstaedter CE, Zhao C, Mattei L, Tanes C, Clarke E, Lauder A, Sherrill-Mix S, Chehoud C, Kelsen J, Conrad M, Collman RG, Baldassano R, Bushman FD, Bittinger K. 2017. Optimizing methods and dodging pitfalls in microbiome research. *Microbiome* **5(1)**:52  
DOI [10.1186/s40168-017-0267-5](https://doi.org/10.1186/s40168-017-0267-5).
- Lauber CL, Zhou N, Gordon JI, Knight R, Fierer N. 2010. Effect of storage conditions on the assessment of bacterial community structure in soil and human-associated samples. *FEMS Microbiology Letters* **307(1)**:80–86 DOI [10.1111/j.1574-6968.2010.01965.x](https://doi.org/10.1111/j.1574-6968.2010.01965.x).
- Lahti L, Sudarshan S. 2017. Tools for microbiome analysis in R. Available at <http://microbiome.github.com/microbiome>.
- Lozupone C, Knight R. 2005. UniFrac: a new phylogenetic method for comparing microbial communities. *Applied and Environmental Microbiology* **71(12)**:8228–8235  
DOI [10.1128/AEM.71.12.8228-8235.2005](https://doi.org/10.1128/AEM.71.12.8228-8235.2005).
- Luz Calle M. 2019. Statistical analysis of metagenomics data. *Genomics & Informatics* **17(1)**:e6  
DOI [10.5808/GI.2019.17.1.e6](https://doi.org/10.5808/GI.2019.17.1.e6).
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal* **17(1)**:10 DOI [10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200).
- McDonald D, Hyde E, Debelius JW, Morton JT, Gonzalez A, Ackermann G, Aksenov AA, Behsaz B, Brennan C, Chen Y, DeRight Goldasich L, Dorrestein PC, Dunn RR, Fahimipour AK, Gaffney J, Gilbert JA, Gogul G, Green JL, Hugenholtz P, Humphrey G, Huttenhower C, Jackson MA, Janssen S, Jeste DV, Jiang L, Kelley ST, Knights D, Kosciolk T, Ladau J, Leach J, Marotz C, Meleshko D, Melnik AV, Metcalf JL, Mohimani H, Montassier E, Navas-Molina J, Nguyen TT, Peddada S, Pevzner P, Pollard KS, Rahnavard G, Robbins-Pianka A, Sangwan N, Shorenstein J, Smarr L, Song SJ, Spector T, Swafford AD, Thackray VG, Thompson LR, Tripathi A, Vázquez-Baeza Y, Vrbanc A, Wischmeyer P, Wolfe E, Zhu Q, Knight R, Greene CS, Mann AE, Amir A, Frazier A, Martino C, Lebrilla C, Lozupone C, Lewis CM Jr, Raison C, Zhang C, Lauber CL, Warinner C, Lowry CA, Callewaert C, Bloss C, Willner D, Galzerani DD, Gonzalez DJ, Mills DA, Chopra D, Gevers D, Berg-Lyons D, Sears DD, Wendel D, Lovelace E, Pierce E, TerAvest E, Bolyen E, Bushman FD, Wu GD, Church GM, Saxe G, Holscher HD, Ugrina I, German JB, Caporaso JG, Wozniak JM, Kerr J, Ravel J, Lewis JD, Suchodolski JS, Jansson JK,

- Hampton-Marcell JT, Bobe J, Raes J, Chase JH, Eisen JA, Monk J, Clemente JC, Petrosino J, Goodrich J, Gauglitz J, Jacobs J, Zengler K, Swanson KS, Lewis K, Mayer K, Bittinger K, Dillon L, Zaramela LS, Schriml LM, Dominguez-Bello MG, Jankowska MM, Blaser M, Pirrung M, Minson M, Kurisu M, Ajami N, Gottel NR, Chia N, Fierer N, White O, Cani PD, Gajer P, Strandwitz P, Kashyap P, Dutton R, Park RS, Xavier RJ, Mills RH, Krajmalnik-Brown R, Ley R, Owens SM, Klemmer S, Matamoros S, Mirarab S, Moorman S, Holmes S, Schwartz T, Eshoo-Anton TW, Vigers T, Pandey V, Treuren WV, Fang X, Zech Xu Z, Jarmusch A, Geier J, Reeve N, Silva R, Kopylova E, Nguyen D, Sanders K, Salido Benitez RA, Heale AC, Abramson M, Waldispühl J, Butyaev A, Drogaris C, Nazarova E, Ball M, Gunderson B, Knight R. 2018. American gut: an open platform for citizen science microbiome research. *mSystems* 3(3):31–49 DOI 10.1128/msystems.00031-18.
- McMurdie PJ, Holmes S. 2013. Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE* 8(4):e61217 DOI 10.1371/journal.pone.0061217.
- Metzler-Zebeli BU, Lawlor PG, Magowan E, Zebeli Q. 2016. Effect of freezing conditions on fecal bacterial composition in pigs. *Animals* 6(3):18 DOI 10.3390/ani6030018.
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB, Simpson GL, Slymos P, Stevens MHH, Szoecs E, Wagner H. 2019. *vegan: community ecology package*. R package version 2.5-6. Available at <https://CRAN.R-project.org/package=vegan>.
- Panek M, Čipčić Paljetak H, Barešić A, Perić M, Matijašić M, Lojkić I, Vranešić Bender D, Krznarić Ž, Verbanac D. 2018. Methodology challenges in studying human gut microbiota—effects of collection, storage, DNA extraction and next generation sequencing technologies. *Scientific Reports* 8(1):5143 DOI 10.1038/s41598-018-23296-4.
- Regan CE, Medill SA, Poissant J, McLoughlin PD. 2020. Causes and consequences of an unusually male-biased adult sex ratio in an unmanaged feral horse population. *Journal of Animal Ecology* 89(12):2909–2921 DOI 10.1111/1365-2656.13349.
- Schliep KP. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics* 27(4):592–593 DOI 10.1093/bioinformatics/btq706.
- Shaw AG, Sim K, Powell E, Cornwell E, Cramer T, McClure ZE, Li MS, Kroll JS. 2016. Latitude in sample handling and storage for infant faecal microbiota studies: the elephant in the room? *Microbiome* 4(1):40 DOI 10.1186/s40168-016-0186-x.
- Sinha R, Abu-Ali G, Vogtmann E, Fodor AA, Ren B, Amir A, Schwager E, Crabtree J, Ma S, Abnet CC, Knight R, White O, Huttenhower C. 2017. The microbiome quality control project consortium, assessment of variation in microbial community amplicon sequencing by the Microbiome Quality Control (MBQC) project consortium. *Nature Biotechnology* 35:1077–1086 DOI 10.1038/nbt.3981.
- Song SJ, Amir A, Metcalf JL, Amato KR, Xu ZZ, Humphrey G, Knight R. 2016. Preservation methods differ in fecal microbiome stability, affecting suitability for field studies. *mSystems* 1(3):e00021-16 DOI 10.1128/mSystems.00021-16.
- Tap J, Cools-Portier S, Pavan S, Druense A, Öhman L, Törnblom H, Simren M, Derrien M. 2019. The effects of the long-term storage of human fecal microbiota samples collected in RINAlater. *Scientific Reports* 9(1):601 DOI 10.1038/s41598-018-36953-5.
- Tedjo DI, Jonkers DMAE, Savelkoul PH, Masclee AA, Van Best N, Pierik MJ, Penders J. 2015. The effect of sampling and storage on the fecal microbiota composition in healthy and diseased subjects. *PLOS ONE* 10(5):e0126685 DOI 10.1371/journal.pone.0126685.

- Vandeputte D, Tito RY, Vanleeuwen R, Falony G, Raes J. 2017.** Practical considerations for large-scale gut microbiome studies. *FEMS Microbiology Reviews* **41(1)**:S154–S167 DOI [10.1093/femsre/fux027](https://doi.org/10.1093/femsre/fux027).
- Vogtmann E, Chen J, Kibriya MG, Chen Y. 2017.** Comparison of fecal collection methods for microbiota studies in Bangladesh. *Applied and Environmental Microbiology* **83(10)**:e00361-17 DOI [10.1128/AEM.00361-17](https://doi.org/10.1128/AEM.00361-17).
- Wang Z, Zolnik CP, Qiu Y, Usyk M, Wang T, Strickler HD, Isasi CR, Kaplan RC, Kurland IJ, Qi Q, Burk RD. 2018.** Comparison of fecal collection methods for microbiome and metabolomics studies. *Frontiers in Cellular and Infection Microbiology* **8**:301 DOI [10.3389/fcimb.2018.00301](https://doi.org/10.3389/fcimb.2018.00301).
- Wright ES. 2015.** DECIPHER: harnessing local sequence context to improve protein multiple sequence alignment. *BMC Bioinformatics* **16(1)**:322 DOI [10.1186/s12859-015-0749-z](https://doi.org/10.1186/s12859-015-0749-z).